

CONCENTRATES 20% FOR BREEDERS -NEW-
REARING + PULLET + LAYING OF HATCHING EGGS
COMPOSITION COMPLETE ANIMAL FEED



| BROILER BREEDERS & LAYER BREEDERS | | REARING | PULLET | LAYING |
|-----------------------------------|---------|-------------------------|-------------------------|-------------------------|
| | | (ca. 0-6/8 weeks) | (ca. 6/8 - 20 weeks) | (20 weeks - end) |
| COMPOSITION OF COMPOUND FEED | | | | |
| | | <i>inclusion</i> 20% | <i>inclusion</i> 20% | <i>inclusion</i> 20% |
| Intraco REARING 20-47-1 | % | 20 | 0 | 0 |
| Intraco LAYER 20-44-1 | % | 0 | 20 | 0 |
| Intraco BREEDER 20-40-1 | % | 0 | 0 | 20 |
| Maize | % | 57 | 58,5 | 59 |
| Wheat pollards/bran | % | 12 | 20 | 15 |
| Limestone | % | 1,25 | 1,5 | 6 |
| Soyabean meal 44% CP | % | 9,75 | / | / |
| Total | % | <u>100</u> | <u>100</u> | <u>100</u> |
| ANALYSIS | | | | |
| Crude protein | % | 19,96 | 16,40 | 14,90 |
| Crude fat | % | 3,63 | 3,80 | 3,63 |
| Crude fiber | % | 3,92 | 4,10 | 3,60 |
| Crude ash | % | 5,36 | 5,29 | 11,13 |
| Calcium | % | 0,87 | 0,92 | 3,31 |
| Total phosphorus | % | 0,62 | 0,63 | 0,56 |
| Phosphorus available Poultry | % | 0,41 | 0,41 | 0,38 |
| Phosphorus digestible Poultry | % | 0,39 | 0,39 | 0,36 |
| Sodium | % | 0,14 | 0,14 | 0,14 |
| Chlorine | % | 0,25 | 0,23 | 0,22 |
| Total Lysine | % | 1,17 | 0,88 | 0,80 |
| Lysine digestible Poultry | % | 0,95 | 0,70 | 0,63 |
| Total Methionine | % | 0,46 | 0,42 | 0,42 |
| Methionine digestible Poultry | % | 0,43 | 0,40 | 0,40 |
| Total Methionine + Cystine | % | 0,77 | 0,70 | 0,67 |
| Meth. + Cyst. digestible Poultry | % | 0,67 | 0,61 | 0,59 |
| Total Tryptophane | % | 0,23 | 0,19 | 0,16 |
| Total Threonine | % | 0,75 | 0,61 | 0,56 |
| Linoleic Acid (C18:2) | % | 1,64 | 1,72 | 1,66 |
| Available Xantophylls | ppm | 10,34 | 10,61 | 10,70 |
| Metabolisable Energy Poultry | Kcal/kg | 2800 | 2850 | / |
| Metabolisable Energy Laying hens | Kcal/kg | / | / | 2825 |