INTRACO Ltd.
Quality ingredients
Expert advice and Responsible Solutions

CONCENTRATES 30% FOR BREEDERS -NEW-

REARING + PULLET + LAYING OF HATCHING EGGS

COMPOSITION COMPLETE ANIMAL FEED

BROILER BREEDERS & LAYER BREEDERS		REARING (ca. 0-6/8 weeks)		PULLET (ca. 6/8 - 20 weeks)		LAYING (20 weeks - end)
COMPOSITION OF COMPOUND FEED						
		high inclusion ***32,5%***	***normal inclusion 30%***	*high inclusion* ***32,5%***	***normal inclusion 30%***	***normal inclusion 30%***
Intraco Broiler 30-44-1	%	32,5	30	0	0	0
Intraco Pullets 30-35-1	%	0	0	32,5	30	0
Intraco Breeder 30-36-1	%	0	0	0	0	30
Maize	%	67,5	57	67,5	63	65
Wheat Pollards/bran	%	0	13	0	7	5
Total	%	**100**	**100**	**100**	**100**	**100**
ANALYSIS						
Crude protein	%	20,10	20,05	17,10	16,92	17,10
Crude fat	%	3,60	3,60	3,75	3,74	3,45
Crude fiber	%	2,95	3,80	3,93	4,27	3,00
Crude ash	%	5,72	5,90	5,92	5,83	10,46
Calcium	%	1,06	1,03	1,10	1,07	3,05
Total phosphorus	%	0,61	0,67	0,60	0,63	0,55
Phosphorus available Poultry	%	0,47	0,45	0,41	0,39	0,40
Phosphorus digestible Poultry	%	0,44	0,42	0,38	0,37	0,35
Sodium	%	0,15	0,15	0,15	0,15	0,15
Total Lysine	%	1,18	1,16	0,83	0,81	0,98
Lysine digestible Poultry	%	0,96	0,94	0,68	0,66	0,78
Total Methionine	%	0,53	0,52	0,34	0,33	0,33
Methionine digestible Poultry	%	0,49	0,47	0,31	0,30	0,30
Total Methionine + Cystine	%	0,83	0,82	0,63	0,63	0,59
Meth. + Cyst. digestible Poultry	%	0,73	0,71	0,53	0,53	0,50
Total Tryptophane	%	0,22	0,23	0,19	0,19	0,19
Total Threonine	%	0,78	0,76	0,66	0,65	0,65
Linoleic Acid (C18:2)	%	1,68	1,63	1,77	1,75	1,61
Available Xantophylls	ppm	12,25	10,35	12,25	11,45	11,80
Metabolisable Energy Poultry	Kcal/kg	2900	2750	2850	2750	/
Metabolisable Energy Laying hens	Kcal/kg	/	/	/	/	2800