

BROILER CONCENTRATE 30% -NEW-
STARTER + GROWER + FINISHER
COMPOSITION COMPLETE ANIMAL FEED



		STARTER (ca. 0-12 days)		GROWER (ca. 12-35 days)		FINISHER (ca. 35 days - slaughter)
COMPOSITION OF COMPOUND FEED						
		<i>high inclusion</i> 32,5%	<i>normal inclusion</i> 30%	<i>high inclusion</i> 32,5%	<i>normal inclusion</i> 30%	<i>normal inclusion</i> 30%
Intraco Broiler 30-44-1	%	32,5	30	32,5	30	30
Maize	%	67,5	50	67,5	57	64
Wheat Pollards/bran	%	0	20	0	13	6
Total	%	<u>100</u>	<u>100</u>	<u>100</u>	<u>100</u>	<u>100</u>
ANALYSIS						
Crude protein	%	20,10	20,50	20,04	20,05	19,60
Crude fat	%	3,60	3,52	3,58	3,60	3,60
Crude fiber	%	2,95	4,21	2,95	3,80	3,30
Crude ash	%	5,72	6,12	5,72	5,90	5,60
Calcium	%	1,06	1,00	1,07	1,00	1,00
Total phosphorus	%	0,61	0,72	0,62	0,67	0,63
Phosphorus available Poultry	%	0,47	0,45	0,46	0,44	0,44
Phosphorus digestible Poultry	%	0,44	0,43	0,42	0,41	0,40
Sodium	%	0,14	0,14	0,14	0,14	0,14
Total Lysine	%	1,18	1,20	1,18	1,16	1,15
Lysine digestible Poultry	%	0,96	0,95	0,96	0,94	0,92
Total Methionine	%	0,53	0,52	0,53	0,51	0,51
Methionine digestible Poultry	%	0,49	0,47	0,50	0,47	0,47
Total Methionine + Cystine	%	0,83	0,83	0,83	0,82	0,80
Meth. + Cyst. digestible Poultry	%	0,73	0,72	0,73	0,72	0,70
Total Tryptophane	%	0,22	0,24	0,22	0,23	0,22
Total Threonine	%	0,78	0,78	0,77	0,76	0,74
Linoleic Acid (C18:2)	%	1,68	1,60	1,68	1,63	1,67
Available Xantophylls	ppm	12,25	9,10	12,20	10,35	11,60
Metabolisable Energy Broilers	Kcal/kg	2750	2400	2750	2550	2700